

Lamotrigine Extended-Release Tablets

Type of Posting	Revision Bulletin
Posting Date	27–Apr–2018
Official Date	01–May–2018
Expert Committee	Chemical Medicines Monographs 4
Reason for Revision	Compliance

In accordance with the Rules and Procedures of the 2015–2020 Council of Experts, the Chemical Medicines Monographs 4 Expert Committee has revised the Lamotrigine Extended-Release Tablets monograph. The purpose for the revision is to add *Dissolution Test 5* to accommodate drug products that were approved with different dissolution conditions and acceptance criteria. The revision necessitates a change in the table numbering in the *Organic Impurities* section.

- *Dissolution Test 5* was validated using an Xterra RP18 brand of L1 column. The typical retention time for lamotrigine is about 2.5 min.

The Lamotrigine Extended-Release Tablets Revision Bulletin supersedes the currently official Lamotrigine Extended-Release Tablets monograph. The Revision Bulletin will be incorporated in *USP 42–NF 37*.

Should you have any questions, please contact Ren-Hwa Yeh, Senior Scientific Liaison (301-998-6818 or rhy@usp.org).

Lamotrigine Extended-Release Tablets

DEFINITION

Lamotrigine Extended-Release Tablets contain NLT 90.0% and NMT 110.0% of the labeled amount of lamotrigine (C₉H₇Cl₂N₃).

IDENTIFICATION

- **A.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*.

ASSAY

• PROCEDURE

Mobile phase: Acetonitrile, water, and trifluoroacetic acid (25:75:0.05)

Diluent: Acetonitrile, methanol, and water (10:20:70)

Standard solution: 0.25 mg/mL of USP Lamotrigine RS in *Diluent*. Sonication may be used to aid dissolution.

Sample stock solution: 1.0–3.0 mg/mL of lamotrigine prepared as follows. Transfer NLT 5 Tablets to a suitable volumetric flask containing 10% of the flask volume of acetonitrile. Allow the Tablets to disperse. Add 20% of the flask volume of methanol. Sonicate for 10 min. Add 30% of the flask volume of 0.1 M hydrochloric acid. Sonicate for 25 min or until a fine, even dispersion is obtained. Allow to cool to room temperature. Dilute with 0.1 M hydrochloric acid to volume. Pass a portion of the solution through a nylon filter of 0.45-µm pore size and use the filtrate.

Sample solution: Nominally 0.2–0.3 mg/mL of lamotrigine in 0.1 M hydrochloric acid from a suitable volume of *Sample stock solution*

Chromatographic system

(See *Chromatography* <621>, *System Suitability*.)

Mode: LC

Detector: UV 270 nm

Column: 4.6-mm × 15-cm; 3-µm packing L1

Column temperature: 40°

Flow rate: 1 mL/min

Injection volume: 5 µL

Run time: NLT 8 times the retention time of lamotrigine

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 1.5%

Analysis

Samples: *Standard solution* and *Sample solution*
Calculate the percentage of the labeled amount of lamotrigine (C₉H₇Cl₂N₃) in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

- r_U = peak response from the *Sample solution*
 r_S = peak response from the *Standard solution*
 C_S = concentration of USP Lamotrigine RS in the *Standard solution* (mg/mL)
 C_U = nominal concentration of lamotrigine in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

Change to read:

• DISSOLUTION <711>

Test 1

Medium 1: 0.01 M hydrochloric acid; 700 mL

Medium 2: 7.8 g of tribasic sodium phosphate, and 22.5 g of sodium dodecyl sulfate in 1 L of water. This solution has a pH of about 12.

Apparatus 2: 50 rpm with sinkers (see *Dissolution* <711>, *Figure 2a*)

Times

For Tablets labeled to contain 25 or 50 mg: 2, 7, 15 h

For Tablets labeled to contain 100, 200, or 250 mg: 2, 5, 12 h

For Tablets labeled to contain 300 mg: 2, 6, 13 h

Procedure: Run the test with *Medium 1* for 2 h. Add 200 mL of *Medium 2*, preheated at 37°. Within 5 min of the addition of *Medium 2*, withdraw the sample for the 2-h time point. Continue the testing by drawing samples at the time points specified in *Table 1*, *Table 2*, or *Table 3*, depending on the label claim.

Diluent: *Medium 1* and *Medium 2* (70:20)

Standard solution: (L/900) mg/mL of USP Lamotrigine RS in *Diluent*, where L is the label claim in mg/Tablet

Sample solution: Pass a suitable portion of the solution under test through a suitable filter of 0.45-µm pore size. Dilute with *Diluent* if necessary.

Blank: *Diluent*

Instrumental conditions

Mode: UV

Analytical wavelength: 260 nm. [NOTE—Depending on the label claim, cells with suitable path lengths may be used.]

Analysis

Samples: *Standard solution* and *Sample solution*
Calculate the percentage of the labeled amount (Q_i) of lamotrigine (C₉H₇Cl₂N₃) dissolved at each time point *i*:

$$\text{Result} = (A_U/A_S) \times C_S \times V \times D \times (1/L) \times 100$$

- A_U = absorbance of the *Sample solution*
 A_S = absorbance of the *Standard solution*
 C_S = concentration of the *Standard solution* (mg/mL)
 V = volume of *Medium*, 900 mL
 D = dilution factor if needed
 L = label claim (mg/Tablet)

Tolerances

For Tablets with 25- or 50-mg label claim: See *Table 1*.

For Tablets with 100-, 200-, or 250-mg label claim: See *Table 2*.

For Tablets with 300-mg label claim: See *Table 3*.

Table 1

Time Point (t)	Time (h)	Amount Dissolved
1	2	NMT 10%
2	7	35%–55%
3	15	NLT 80%

Table 2

Time Point (i)	Time (h)	Amount Dissolved	
		100 mg, 200 mg	250 mg
1	2	NMT 10%	NMT 10%
2	5	20%–45%	20%–40%
3	12	NLT 80%	NLT 80%

Table 3

Time Point (i)	Time (h)	Amount Dissolved
1	2	NMT 10%
2	6	25%–45%
3	13	NLT 80%

The percentages of the labeled amount of lamotrigine ($C_9H_7Cl_2N_3$) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*.

Test 2: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 2*.

Acid stage medium: 0.01 M hydrochloric acid; 700 mL

Buffer stage stock medium: 2.83 g of sodium phosphate monobasic, 1.72 g of sodium hydroxide, and 22.5 g of sodium dodecyl sulfate in 1 L of water


Buffer stage medium: *Acid stage medium* and *Buffer stage stock medium* (70:20); 900 mL. Adjust with solution A (phosphoric acid in water prepared by diluting 1 mL of phosphoric acid with water to 50 mL) or 0.1 N sodium hydroxide, if necessary, to a pH of 6.8. Record the required volume of solution A or 0.1 N sodium hydroxide for adjustment of pH to 6.8.

Apparatus 2: 50 rpm with stationary tablet basket. See *Figures 1* and *2*.

Times

For Tablets labeled to contain 25 or 50 mg: 2 h in *Acid stage medium*; 4, 7, 9, and 15 h in *Buffer stage medium*

For Tablets labeled to contain 100, 200, or 300 mg: 2 h in *Acid stage medium*; 3, 5, 7, and 12 h in *Buffer stage medium*

**NOTES**

1. Rod and Basket with a Tablet cover placed in the horizontal diagonal of the basket.
2. Basket and Tablet cover material; stainless steel.
3. Basket gauze wire size: 8 mesh.

Figure 1. Stationary tablet basket.




Figure 2. Drug release stationary tablet basket configuration diagram.

[NOTE—The times in the *Buffer stage medium* include the time in the *Acid stage medium*.]

Procedure: Run the test with *Acid stage medium* for 2 h followed by collecting the *Acid stage medium* sample and replacing it with the same volume of *Acid stage medium*. Add 200 mL of *Buffer stage stock medium* to the above solution. If necessary, add either solution A or 0.1 N sodium hydroxide to the solution to reach a pH of 6.8. Continue the testing by drawing samples at the time points specified in *Table 4* or *Table 5*, depending on the label claim. Replace each of the volumes withdrawn with an equal volume of *Buffer stage medium*.

Buffer: Dissolve 2.76 g of sodium phosphate monobasic in 1 L of water. Add 2 mL of triethylamine and adjust with solution A to a pH of 7.0.

Mobile phase: Methanol and *Buffer* (55:45)

Standard stock solution: 1.4 mg/mL of USP Lamotrigine RS in methanol

Acid stage standard solution: (L/900) mg/mL of USP Lamotrigine RS from *Standard stock solution*, in *Acid stage medium*, where L is the label claim in mg/Tablet

Buffer stage standard solution: (L/900) mg/mL of USP Lamotrigine RS from *Standard stock solution*, in *Buffer stage medium*, where L is the label claim in mg/Tablet

Acid stage sample solution: Withdraw a 10.0-mL aliquot, and pass a portion of the solution under test through a suitable filter of 0.45- μ m pore size. Replace the 10.0-mL aliquot withdrawn for analysis with a 10.0-mL aliquot of *Acid stage medium*.

Buffer stage sample solution: Withdraw a 10.0-mL aliquot, and pass a portion of the solution under test through a suitable filter of 0.45- μ m pore size. Replace the 10.0-mL aliquot withdrawn for analysis with a 10.0-mL aliquot of *Buffer stage medium*.

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 270 nm

Column: 4.6-mm \times 15-cm; 5- μ m packing L7

Flow rate: 1 mL/min

Injection volume

For 25-mg Tablets: 80 μ L

For 50-mg Tablets: 40 μ L

For 100-mg Tablets: 20 μ L

For 200- or 300-mg Tablets: 10 μ L

Run time: NLT 1.8 times the retention time of lamotrigine

System suitability

Samples: *Acid stage standard solution* and *Buffer stage standard solution*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Acid stage standard solution*, *Acid stage sample solution*, *Buffer stage standard solution*, and *Buffer stage sample solution*

Calculate the percentage (Q_A) of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved in the *Acid stage medium*:

$$\text{Result}_i = (r_U/r_S) \times C_S \times V_A \times (1/L) \times 100$$

r_U = peak response from the *Acid stage sample solution*

r_S = peak response from the *Acid stage standard solution*

C_S = concentration of USP Lamotrigine RS in the *Acid stage standard solution* (mg/mL)

V_A = volume of the *Acid stage medium*, 700 mL

L = label claim (mg/Tablet)

Calculate the concentration (C_i) of lamotrigine ($C_9H_7Cl_2N_5$) in the sample withdrawn from the vessel at each time point (i) during the buffer stage:

$$\text{Result}_i = (r_U/r_S) \times C_S$$

- r_U = peak response from the *Buffer stage sample solution* at time point i
 r_S = peak response from the *Buffer stage standard solution*
 C_S = concentration of USP Lamotrigine RS in the *Buffer stage standard solution* (mg/mL)

Calculate the percentage of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved at each time point (i) during the buffer stage:

$$\text{Result}_1 = [C_1 \times V_B \times (1/L) \times 100] + (Q_A \times V_S/V_A)$$

$$\text{Result}_2 = \{[(C_2 \times V_B) + (C_1 \times V_S)] \times (1/L) \times 100\} + (Q_A \times V_S/V_A)$$

$$\text{Result}_3 = \{[(C_3 \times V_B) + [(C_2 + C_1) \times V_S]] \times (1/L) \times 100\} + (Q_A \times V_S/V_A)$$

$$\text{Result}_4 = \{[(C_4 \times V_B) + [(C_3 + C_2 + C_1) \times V_S]] \times (1/L) \times 100\} + (Q_A \times V_S/V_A)$$

- C_i = concentration of lamotrigine in the *Buffer stage sample solution* withdrawn at time point i (mg/mL)
 V_B = volume of the *Buffer stage medium*, 900 mL
 L = label claim (mg/Tablet)
 Q_A = the percentage of the labeled amount of lamotrigine dissolved in the *Acid stage medium*
 V_S = volume of the sample solution withdrawn at each time point (i) during the acid stage or buffer stage (mL)
 V_A = volume of the *Acid stage medium*, 700 mL

Tolerances

For Tablets labeled to contain 25 or 50 mg: See Table 4.

For Tablets labeled to contain 100, 200, or 300 mg: See Table 5.

Table 4

Time Point (i)	Time (h)	Amount Dissolved
1	2	NMT 10%
2	4	5%–25%
3	7	30%–50%
4	9	50%–70%
5	15	NLT 80%

Table 5

Time Point (i)	Time (h)	Amount Dissolved
1	2	NMT 10%
2	3	5%–20%
3	5	25%–50%
4	7	50%–70%
5	12	NLT 80%

The percentages of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved at the times specified conform to *Dissolution* (711), *Acceptance Table 2*.

Test 3: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 3*.

Acid stage medium: 0.01 M hydrochloric acid; 700 mL

Buffer stage stock medium: 7.8 g of sodium phosphate tribasic and 22.5 g of sodium dodecyl sulfate in 1 L of water

Buffer stage medium: *Acid stage medium* and *Buffer stage stock medium* (70:20); 900 mL. Adjust with 2 N hydrochloric acid TS or 2 N sodium hydroxide TS, if necessary, to a pH of 6.8.

Apparatus 2: 50 rpm with stationary tablet basket. See *Figures 1* and *2* in *Test 2*.

Times

For Tablets labeled to contain 25, 50, 100, or 200 mg: 2 h in *Acid stage medium*; 4, 7, and 14 h in *Buffer stage medium*

[NOTE—The times in the *Buffer stage medium* include the time in the *Acid stage medium*.]

Procedure: Run the test with *Acid stage medium* for 2 h followed by collecting the *Acid stage medium* sample and replacing it with the same volume of *Acid stage medium*. Add 200 mL of *Buffer stage stock medium* to the above solution. Continue the testing by drawing samples at the time points specified in *Table 6*. Replace each of the volumes withdrawn with an equal volume of the *Buffer stage medium*.

Buffer: Dissolve 2.72 g of potassium phosphate monobasic in 1 L of water and adjust with dilute phosphoric acid to a pH of 3.7.

Mobile phase: Methanol, acetonitrile, and *Buffer* (50:15:35)

Standard stock solution: 0.6 mg/mL of USP Lamotrigine RS in methanol. Sonicate to dissolve as needed.

Acid stage standard solution: 0.036 mg/mL of USP Lamotrigine RS from *Standard stock solution*, in *Acid stage medium*

Buffer stage standard solution: ($L/900$) mg/mL of USP Lamotrigine RS from *Standard stock solution*, in *Buffer stage medium*, where L is the label claim in mg/Tablet

Acid stage sample solution: Withdraw a 10.0-mL aliquot and pass a portion of the solution under test through a suitable filter of 0.45- μ m pore size, discarding the first 5 mL of the filtrate. Replace the 10.0-mL aliquot withdrawn for analysis with a 10.0-mL aliquot of *Acid stage medium*.

Buffer stage sample solution: Withdraw a 10.0-mL aliquot and pass a portion of the solution under test through a suitable filter of 0.45- μ m pore size. Replace the 10.0-mL aliquot withdrawn for analysis with a 10.0-mL aliquot of *Buffer stage medium*.

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 270 nm

Column: 4.6-mm \times 25-cm; 5- μ m packing L1

Column temperature: 35°

Flow rate: 1 mL/min

Injection volume: 10 μ L

Run time: NLT 1.7 times the retention time of lamotrigine

System suitability

Samples: *Acid stage standard solution* and *Buffer stage standard solution*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Acid stage standard solution, Acid stage sample solution, Buffer stage standard solution, and Buffer stage sample solution*

Calculate the percentage (Q_A) of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved in the *Acid stage medium*:

$$\text{Result}_i = (r_U/r_S) \times C_S \times V_A \times (1/L) \times 100$$

- r_U = peak response from the *Acid stage sample solution*
- r_S = peak response from the *Acid stage standard solution*
- C_S = concentration of USP Lamotrigine RS in the *Acid stage standard solution* (mg/mL)
- V_A = volume of the *Acid stage medium*, 700 mL
- L = label claim (mg/Tablet)

Calculate the concentration (C_i) of lamotrigine ($C_9H_7Cl_2N_5$) in the sample withdrawn from the vessel at each time point (i) during the buffer stage:

$$\text{Result}_i = (r_U/r_S) \times C_S$$

- r_U = peak response from the *Buffer stage sample solution* at time point i
- r_S = peak response from the *Buffer stage standard solution*
- C_S = concentration of USP Lamotrigine RS in the *Buffer stage standard solution* (mg/mL)

Calculate the percentage of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved at each time point (i) during the buffer stage:

$$\begin{aligned} \text{Result}_1 &= [C_1 \times V_B \times (1/L) \times 100] + (Q_A \times V_S/V_A) \\ \text{Result}_2 &= \{[(C_2 \times V_B) + (C_1 \times V_S)] \times (1/L) \times 100\} + (Q_A \times V_S/V_A) \\ \text{Result}_3 &= \{[(C_3 \times V_B) + [(C_2 + C_1) \times V_S]] \times (1/L) \times 100\} + (Q_A \times V_S/V_A) \end{aligned}$$

- C_i = concentration of lamotrigine in the *Buffer stage sample solution* withdrawn at time point i (mg/mL)
- V_B = volume of the *Buffer stage medium*, 900 mL
- L = label claim (mg/Tablet)
- Q_A = the percentage of the labeled amount of lamotrigine dissolved in the *Acid stage medium*
- V_S = volume of the sample solution withdrawn at each time point (i) during the acid stage or buffer stage (mL)
- V_A = volume of the *Acid stage medium*, 700 mL

Tolerances

For Tablets labeled to contain 25, 50, 100, or 200 mg: See *Table 6*.

Table 6

Time Point (i)	Time (h)	Amount Dissolved (%)
1	2	NMT 10
2	4	NMT 25
3	7	36–61
4	14	NLT 85

The percentages of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved at the times specified conform to *Dissolution <711>*, *Acceptance Table 2*.

Test 4: If the product complies with this test, the labeling indicates that it meets *USP Dissolution Test 4*.

Acid stage medium: 0.01 M hydrochloric acid; 700 mL

Buffer stage stock medium: 7.8 g of sodium phosphate tribasic and 22.5 g of sodium dodecyl sulfate in 1 L of water

Buffer stage medium: *Acid stage medium* and *Buffer stage stock medium* (70:20); 900 mL

Apparatus 2: 50 rpm with sinkers

Times

For Tablets labeled to contain 25, 50, 100, 200, or 300 mg: 2 h in *Acid stage medium*; 9 and 17 h in *Buffer stage medium*

[NOTE—The times in the *Buffer stage medium* include the time in the *Acid stage medium*.]

Procedure: Run the test with *Acid stage medium* for 2 h followed by collecting the *Acid stage medium* sample and replacing it with the same volume of *Acid stage medium*. Add 200 mL of *Buffer stage stock medium* to the above solution. Continue the testing by drawing samples at the time points specified in *Table 7*. Replace each of the volumes withdrawn with an equal volume of the *Buffer stage medium*.

Buffer: Dissolve 4.1 g of potassium phosphate monobasic in 900 mL of water and adjust with dilute phosphoric acid to a pH of 2.0, and then dilute with water to 1 L. Add 1.25 g of sodium 1-hexanesulfonate to the solution and mix.

Mobile phase: Acetonitrile and *Buffer* (25:75)

Acid stage standard stock solution 1: 0.07 mg/mL of USP Lamotrigine RS prepared as follows. Transfer an appropriate amount of USP Lamotrigine RS to a suitable volumetric flask and add 20% of the flask volume of methanol. Sonicate to dissolve and dilute with *Acid stage medium* to volume. Further dilute this solution with *Acid stage medium* to obtain a final concentration of 0.07 mg/mL.

Acid stage standard stock solution 2: 0.14 mg/mL of USP Lamotrigine RS prepared as follows. Transfer an appropriate amount of USP Lamotrigine RS to a suitable volumetric flask and add 20% of the flask volume of methanol. Sonicate to dissolve and dilute with *Acid stage medium* to volume. Further dilute this solution with *Acid stage medium* to obtain a final concentration of 0.14 mg/mL.

Acid stage standard solution: $0.1 \times (L/700)$ mg/mL of USP Lamotrigine RS either from *Acid stage standard stock solution 1* for Tablets labeled to contain 25, 50, 100, and 200 mg, or from *Acid stage standard stock solution 2* for Tablets labeled to contain 300 mg, in *Acid stage medium*, where L is the label claim in mg/Tablet. Pass a portion of the solution through a suitable filter of 0.45- μ m pore size, discarding the first 2–3 mL of the filtrate.

Buffer stage standard stock solution 1: 0.55 mg/mL of USP Lamotrigine RS prepared as follows. Transfer an appropriate amount of USP Lamotrigine RS to a suitable volumetric flask and add 10% of the flask volume of methanol. Sonicate to dissolve and dilute with *Buffer stage medium* to volume.

Buffer stage standard stock solution 2: 1.1 mg/mL of USP Lamotrigine RS prepared as follows. Transfer an appropriate amount of USP Lamotrigine RS to a suitable volumetric flask and add 20% of the flask

volume of methanol. Sonicate to dissolve and dilute with *Buffer stage medium* to volume.

Buffer stage standard solution: ($L/900$) mg/mL of USP Lamotrigine RS either from *Buffer stage standard stock solution 1* for Tablets labeled to contain 25, 50, 100, and 200 mg, or from *Buffer stage standard stock solution 2* for Tablets labeled to contain 300 mg, in *Buffer stage medium*, where L is the label claim in mg/Tablet. Pass a portion of the solution through a suitable filter of 0.45- μ m pore size, discarding the first 2–3 mL of the filtrate.

Acid stage sample solution: Withdraw a 10.0-mL aliquot and pass a portion of the solution under test through a suitable filter of 0.45- μ m pore size, discarding the first 2–3 mL of the filtrate. Replace the 10.0-mL aliquot withdrawn for analysis with a 10.0-mL aliquot of *Acid stage medium*. Pass a portion of the solution through a suitable filter of 0.45- μ m pore size, discarding the first 2–3 mL of the filtrate.

Buffer stage sample solution: Withdraw a 10.0-mL aliquot and pass a portion of the solution under test through a suitable filter of 0.45- μ m pore size. Replace the 10.0-mL aliquot withdrawn for analysis with a 10.0-mL aliquot of *Buffer stage medium*. Pass a portion of the solution through a suitable filter of 0.45- μ m pore size, discarding the first 2–3 mL of the filtrate.

Chromatographic system

(See *Chromatography* (621), *System Suitability*.)

Mode: LC

Detector: UV 270 nm

Column: 4.6-mm \times 25-cm; 5- μ m packing L7

Column temperature: 60°

Flow rate: 1 mL/min

Injection volume: 10 μ L

Run time: NLT 2 times the retention time of lamotrigine

System suitability

Samples: *Acid stage standard solution* and *Buffer stage standard solution*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Acid stage standard solution*, *Acid stage sample solution*, *Buffer stage standard solution*, and *Buffer stage sample solution*

Calculate the percentage (Q_A) of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved in the *Acid stage medium*:

$$\text{Result}_i = (r_U/r_S) \times C_S \times V_A \times (1/L) \times 100$$

r_U = peak response from the *Acid stage sample solution*

r_S = peak response from the *Acid stage standard solution*

C_S = concentration of USP Lamotrigine RS in the *Acid stage standard solution* (mg/mL)

V_A = volume of the *Acid stage medium*, 700 mL

L = label claim (mg/Tablet)

Calculate the concentration (C_i) of lamotrigine ($C_9H_7Cl_2N_5$) in the sample withdrawn from the vessel at each time point (i) during the buffer stage:

$$\text{Result}_i = (r_U/r_S) \times C_S$$

r_U = peak response from the *Buffer stage sample solution* at time point i

r_S = peak response from the *Buffer stage standard solution*

C_S = concentration of USP Lamotrigine RS in the *Buffer stage standard solution* (mg/mL)

Calculate the percentage of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved at each time point (i) during the buffer stage:

$$\text{Result}_1 = [C_1 \times V_B \times (1/L) \times 100] + (Q_A \times V_S/V_A)$$

$$\text{Result}_2 = \{[(C_2 \times V_B) + (C_1 \times V_S)] \times (1/L) \times 100\} + (Q_A \times V_S/V_A)$$

C_i = concentration of lamotrigine in the *Buffer stage sample solution* withdrawn at time point i (mg/mL)

V_B = volume of the *Buffer stage medium*, 900 mL

L = label claim (mg/Tablet)

Q_A = the percentage of the labeled amount of lamotrigine dissolved in the *Acid stage medium*

V_S = volume of the sample solution withdrawn at each time point (i) during the acid stage or buffer stage (mL)

V_A = volume of the *Acid stage medium*, 700 mL

Tolerances

For Tablets labeled to contain 25, 50, 100, 200, or 300 mg: See *Table 7*.

Table 7

Time Point (i)	Time (h)	Amount Dissolved (%)
1	2	NMT 10
2	9	35–55
3	17	NLT 80

The percentages of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved at the times specified conform to *Dissolution* (711), *Acceptance Table 2*. (RB 1-Nov-2017)

Test 5: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 5*.

Acid stage medium: 0.01 M hydrochloric acid; 710 mL

Buffer stage stock medium: 3.36 g of anhydrous tribasic sodium phosphate and 22.5 g of sodium dodecyl sulfate in 1 L of water

Buffer stage medium: *Acid stage medium* and *Buffer stage stock medium* (70:20); 900 mL. Adjust with hydrochloric acid or 5 N sodium hydroxide TS, if necessary, to a pH of 6.8.

Apparatus 2: 50 rpm with sinkers

Times

For Tablets labeled to contain 25, 50, 100, or 200 mg: 2 h in *Acid stage medium*; 4, 7, and 17 h in *Buffer stage medium*

[NOTE—The times in the *Buffer stage medium* include the time in *Acid stage medium*.]

Procedure: Run the test with *Acid stage medium* for 2 h followed by collecting the *Acid stage medium* sample. Add 200 mL of *Buffer stage stock medium* to this solution. Continue the testing by drawing samples at the time points specified in *Table 8*.

Buffer: Dissolve 3.45 g of monobasic sodium phosphate in 1 L of water, and adjust with

phosphoric acid to a pH of 3.3. To this solution, add 5.77 g of sodium dodecyl sulfate, and mix well.

Mobile phase: Acetonitrile and Buffer (45:55)

Standard solution: 0.22 mg/mL of USP Lamotrigine RS in *Mobile phase*. Sonicate to dissolve as needed.

Acid stage sample solution: Withdraw a 10.0-mL aliquot, and pass a portion of the solution under test through a suitable full flow filter of 10- μ m pore size.

Buffer stage sample solution: Withdraw a 2.5-mL aliquot, and pass a portion of the solution under test through a suitable full flow filter of 10- μ m pore size.

Chromatographic system

(See *Chromatography* <621>, *System Suitability*.)

Mode: LC

Detector: UV 266 nm

Column: 4.6-mm \times 5-cm; 3.5- μ m packing L1

Column temperature: 30°

Flow rate: 1 mL/min

Injection volume: 10 μ L

Run time: NLT 1.6 times the retention time of lamotrigine

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 1.5

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution*, *Acid stage sample solution*, and *Buffer stage sample solution*

Calculate the percentage (Q_A) of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved in the *Acid stage medium*:

$$\text{Result}_i = (r_U/r_S) \times C_S \times V_A \times (1/L) \times 100$$

r_U = peak response from the *Acid stage sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of USP Lamotrigine RS in the *Standard solution* (mg/mL)

V_A = volume of the *Acid stage medium*, 710 mL

L = label claim (mg/Tablet)

Calculate the concentration (C_i) of lamotrigine ($C_9H_7Cl_2N_5$) in the sample withdrawn from the vessel at each time point (i) during the buffer stage:

$$\text{Result}_i = (r_U/r_S) \times C_S$$

r_U = peak response from the *Buffer stage sample solution* at time point i

r_S = peak response from the *Standard solution*

C_S = concentration of USP Lamotrigine RS in the *Standard solution* (mg/mL)

Calculate the percentage of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved at each time point (i) during the buffer stage:

$$\begin{aligned} \text{Result}_1 &= [C_1 \times V_B \times (1/L) \times 100] + (Q_A \times V_S/V_A) \\ \text{Result}_2 &= \{[C_2 \times (V_B - V_S) + (C_1 \times V_S)] \times (1/L) \times 100\} + \\ &\quad (Q_A \times V_S/V_A) \\ \text{Result}_3 &= \{([C_3 \times (V_B - 2V_S)] + [(C_2 + C_1) \times V_S]) \times (1/L) \times \\ &\quad 100\} + (Q_A \times V_S/V_A) \end{aligned}$$

C_i = concentration of lamotrigine in the *Buffer stage sample solution* withdrawn at time point i (mg/mL)

V_B = volume of the *Buffer stage medium*, 900 mL

L = label claim (mg/Tablet)

Q_A = the percentage of the labeled amount of lamotrigine dissolved in the *Acid stage medium*

V_S = volume of the *Sample solution* withdrawn at each time point (i), 10 mL for acid stage or 2.5 mL for buffer stage

V_A = volume of the *Acid stage medium*, 710 mL

Tolerances

For Tablets labeled to contain 25, 50, 100, or 200 mg: See *Table 8*.

Table 8

Time Point (i)	Time (h)	Amount Dissolved	
		25, 50, and 200 mg/Tablet (%)	100 mg/Tablet (%)
1	2	NMT 10	NMT 10
2	4	10–30	10–30
3	7	35–60	40–65
4	17	NLT 80	NLT 80

The percentages of the labeled amount of lamotrigine ($C_9H_7Cl_2N_5$) dissolved at the times specified conform to *Dissolution* <711>, *Acceptance Table 2*. \blacktriangle (RB 1-May-2018)

- **UNIFORMITY OF DOSAGE UNITS (905):** Meet the requirements

IMPURITIES

Change to read:

- **ORGANIC IMPURITIES**

Mobile phase, Sample solution, and Chromatographic system: Proceed as directed in the *Assay*.

Diluent 1: Acetonitrile, methanol, and 0.1 M hydrochloric acid (10:20:70)

Diluent 2: Acetonitrile, methanol, and water (10:20:70)

System suitability stock solution: 0.025 mg/mL of USP Lamotrigine Related Compound C RS in *Diluent 1*

System suitability solution: 1.25 μ g/mL of USP Lamotrigine Related Compound C RS and 0.25 mg/mL of USP Lamotrigine RS in *Diluent 2* prepared as follows. Transfer a suitable amount of USP Lamotrigine RS to a suitable volumetric flask. Transfer a suitable volume of *System suitability stock solution* to the flask. Dissolve and dilute with *Diluent 2* to volume.

System suitability

Sample: *System suitability solution*

Suitability requirements

Resolution: NLT 10 between the lamotrigine and lamotrigine related compound C peaks

Signal-to-noise ratio: NLT 100 for lamotrigine related compound C

Analysis

Sample: *Sample solution*

Calculate the percentage of each degradation product in the portion of Tablets taken:

$$\text{Result} = (r_U/r_T) \times 100$$

r_U = response of each impurity from the *Sample solution*

8 Lamotrigine

Revision Bulletin
Official May 1, 2018

r_T = sum of all of the impurity peak responses and the lamotrigine peak response from the *Sample solution*

Acceptance criteria: See Table **▲9.▲** (RB 1-May-2018)
Disregard peaks less than 0.05%.

Table ▲9▲ (RB 1-May-2018)

Name	Relative Retention Time	Acceptance Criteria, NMT (%)
Lamotrigine	1.0	—
Lamotrigine related compound C	1.7	0.3
Lamotrigine dimer ^a	6.0	0.2
Any individual unspecified degradation product	—	0.2

Table ▲9▲ (RB 1-May-2018) (continued)

Name	Relative Retention Time	Acceptance Criteria, NMT (%)
Total impurities	—	0.5

^a This is either lamotrigine *o*-dimer [*N*⁵,*N*^{5'}-methylenebis(6-(2,3-dichlorophenyl)-1,2,4-triazine-3,5-diamine)] or lamotrigine *p*-dimer [*N*³,*N*^{3'}-methylenebis(6-(2,3-dichlorophenyl)-1,2,4-triazine-3,5-diamine)].

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in well-closed containers. Store at controlled room temperature.
- **LABELING:** When more than one *Dissolution* test is given, the labeling states the *Dissolution* test used only if *Test 1* is not used.
- **USP REFERENCE STANDARDS (11)**
USP Lamotrigine RS
USP Lamotrigine Related Compound C RS
3-Amino-6-(2,3-dichlorophenyl)-1,2,4-triazin-5(4*H*)-one.
C₉H₆Cl₂N₄O 257.08