

Mesalamine Delayed-Release Tablets

Type of Posting	Revision Bulletin
Posting Date	31–July–2020
Official Date	01–Nov–2020
Expert Committee	Chemical Medicines Monographs 2
Reason for Revision	Compliance

In accordance with the Rules and Procedures of the 2015–2020 Council of Experts, the Chemical Medicines Monographs 2 Expert Committee has revised the Mesalamine Delayed-Release Tablets monograph. The purpose for the revision is to add *Dissolution Test 4* to accommodate FDA-approved drug products with different dissolution conditions and/or tolerances than the existing dissolution tests.

Additionally, minor editorial changes have been made to update the monograph to current *USP* style.

The Mesalamine Delayed-Release Tablets Revision Bulletin replaces the version which is scheduled to become official on Nov. 1, 2020.

Should you have any questions, please contact Tsion Billign, Scientific Liaison (301-816-8286 or tb@usp.org).

Mesalamine Delayed-Release Tablets

DEFINITION

Mesalamine Delayed-Release Tablets contain NLT 90.0% and NMT 110.0% of the labeled amount of mesalamine (C₇H₇NO₃).

IDENTIFICATION

- **A. [SPECTROSCOPIC IDENTIFICATION TESTS \(197\)](#), [Infrared Spectroscopy](#): 197K**

Sample: To about 50 mL of [water](#) add a quantity of finely powdered Tablets, nominally equivalent to about 800 mg of mesalamine. Boil the mixture for about 5 min, with constant stirring. Filter the hot solution, and allow the filtrate to cool. Collect the precipitated crystals, and dry at about 110°.

Acceptance criteria: Meet the requirements

Add the following:

- ▲ ● **B.** The retention time of the mesalamine peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*. ▲ (USP 1-Aug-2020)

ASSAY

Change to read:

- **PROCEDURE**

Mobile phase: Dissolve 4.3 g of [sodium 1-octanesulfonate](#) in 1 L of [water](#). Adjust with [phosphoric acid](#) to a pH of 2.15, pass through a filter of 0.45-µm or finer pore size, and degas.

System suitability stock solution: 0.1 mg/mL each of 3-aminosalicylic acid and [USP Salicylic Acid RS](#) prepared as follows. Transfer about 20 mg each of [3-aminosalicylic acid](#) and [USP Salicylic Acid RS](#) to a 200-mL volumetric flask. Dissolve in 50 mL of 1 N [hydrochloric acid](#), sonicate to dissolve, dilute with [water](#) to volume, and mix.

System suitability solution: 0.01 mg/mL each of [3-aminosalicylic acid](#) and ▲ [USP Salicylic Acid RS](#) ▲ (USP 1-Aug-2020) in water from the *System suitability stock solution*

Standard stock solution: 1 mg/mL of [USP Mesalamine RS](#) prepared as follows. Transfer about 25 mg of [USP Mesalamine RS](#) to a 25-mL volumetric flask. Dissolve in 5 mL of 0.25 N [hydrochloric acid](#), sonicate to dissolve, dilute with [water](#) to volume, and mix.

Standard solution: About 0.2 mg/mL of [USP Mesalamine RS](#) and 0.001 mg/mL of 3-aminosalicylic acid prepared as follows. Transfer 10.0 mL of the *Standard stock solution* and 5.0 mL of the *System suitability solution* to a 50-mL volumetric flask. Dilute with [water](#) to volume.

▲ **Sample stock solution:** Nominally 0.8 mg/mL of mesalamine prepared as follows. Transfer a portion nominally equivalent to about 400 mg of mesalamine, from NLT 20 finely powdered Tablets, to a 500-mL volumetric flask. Add 50 mL of 1 N [hydrochloric acid](#), and sonicate to dissolve. Shake by mechanical means for 10 min, dilute with [water](#) to volume, mix, and pass through a filter of 0.5-µm or finer pore size. ▲ (USP 1-Aug-2020)

Sample solution: ▲ Nominally 0.2 mg/mL of mesalamine prepared by diluting the *Sample stock solution* with [water](#) ▲ (USP 1-Aug-2020)

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 230 nm

▲ (USP 1-Aug-2020)

Column: 4.6-mm × 3.3-cm; 3-μm base-deactivated packing L1

Flow rate: 2 mL/min

Injection volume: 20 μL

System suitability

Sample: *Standard solution*

▲ [NOTE—The relative retention times for salicylic acid, mesalamine, and 3-aminosalicylic acid are about 0.5, 1.0, and 1.6, respectively.] ▲ (USP 1-Aug-2020)

Suitability requirements

Resolution: NLT 2 between mesalamine and salicylic acid; NLT 2 between mesalamine and 3-aminosalicylic acid

Tailing factor: NMT 2

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of mesalamine (C₇H₇NO₃) in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of mesalamine from the *Sample solution*

r_S = peak response of mesalamine from the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of mesalamine in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

Change to read:

- [DISSOLUTION \(711\)](#)

Test 1

Solution A: Transfer about 43.35 g of [monobasic potassium phosphate](#) and 1.65 g of [sodium hydroxide](#) to a 2-L volumetric flask. Dissolve in and dilute with [water](#) to volume, and mix. Adjust with 1 N [sodium hydroxide](#) or [phosphoric acid](#) to a pH of 6.0, and mix.

Solution B: Transfer 133.6 g of [sodium hydroxide](#) to a 2-L volumetric flask, dissolve in and dilute with [water](#) to volume, and mix.

Medium

Acid stage: 500 mL of 0.1 N [hydrochloric acid](#)

Buffer stages: 900 mL of *Solution A*

Apparatus 2

Acid stage: 100 rpm

Buffer stage 1: 100 rpm

Buffer stage 2: 50 rpm

Times

Acid stage: 2 h

Buffer stage 1: 1 h

Buffer stage 2: 90 min

Acid stage: After 2 h of operation, withdraw an aliquot of the fluid, discard the remaining solution, and retain the Tablets in proper order so that each will be returned later to its respective vessel. Blot the Tablets with a paper towel to dry, and proceed immediately as directed in *Buffer stage 1*.

Standard solution: A known concentration of [USP Mesalamine RS](#) in *Medium*, equivalent to about 1% of the labeled amount of mesalamine ($C_7H_7NO_3$)

Sample solution: Filter portions of the solution under test, and suitably dilute with *Medium*, if necessary.

Instrumental conditions

Mode: UV

Analytical wavelength: 302 nm (maximum absorbance)

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved:

$$\text{Result} = (A_U/A_S) \times C_S \times V \times (1/L) \times 100$$

A_U = absorbance of the *Sample solution*

A_S = absorbance of the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 500 mL

L = label claim of mesalamine (mg/Tablet) ▲ (USP 1-Aug-2020)

Tolerances: See [Table 1](#). Continue testing through all levels unless the results conform at an earlier level.

Buffer stage 1: [NOTE—Use *Solution A* that has been equilibrated to a temperature of $37 \pm 0.5^\circ$.] Transfer *Solution A* to each of the dissolution vessels, and place each Tablet from the *Acid stage* into its respective vessel. After 1 h, remove a 50-mL aliquot, and proceed immediately as directed in *Buffer stage 2*.

Standard solution: A known concentration of [USP Mesalamine RS](#) in *Medium*, equivalent to about 1% of the labeled amount of mesalamine ($C_7H_7NO_3$)

Sample solution: Filter portions of the solution under test, and suitably dilute with *Medium*, if necessary.

Instrumental conditions

Mode: UV

Analytical wavelength: 330 nm (maximum absorbance)

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved:

$$\text{Result} = (A_U/A_S) \times C_S \times V \times (1/L) \times 100$$

A_U = absorbance of the *Sample solution*

A_S = absorbance of the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 900 mL

L = label claim of mesalamine (mg/Tablet) [▲] (USP 1-Aug-2020)

Tolerances: See [Table 1](#). Continue testing through all levels unless the results conform at an earlier level.

Table 1

Level	Number Tested	Acceptance Criteria
L_1	6	No individual value exceeds 1% dissolved.
L_2	6	Average of the 12 units ($L_1 + L_2$) is NMT 1% dissolved, and no individual unit is greater than 10% dissolved.
L_3	12	Average of the 24 units ($L_1 + L_2 + L_3$) is NMT 1% dissolved, and NMT 1 individual unit is greater than 10% dissolved.

Buffer stage 2: Add 50 mL of *Solution B* to each dissolution vessel to adjust to a pH of 7.2, and continue the run.

Standard solution: A known concentration of [USP Mesalamine RS](#) in *Medium*

Sample solution: Filter portions of the solution under test, and suitably dilute with *Medium*, if necessary.

Instrumental conditions

Mode: UV

Analytical wavelength: 332 nm (maximum absorbance)

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved:

$$\text{▲Result} = (A_U/A_S) \times C_S \times V \times (1/L) \times 100$$

A_U = absorbance of the *Sample solution*

A_S = absorbance of the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 900 mL

L = label claim of mesalamine (mg/Tablet) [▲] (USP 1-Aug-2020)

Tolerances: NLT 80% (Q) of the labeled amount of mesalamine ($C_7H_7NO_3$) is dissolved. The requirements are met if the quantities dissolved from the product conform to [Dissolution \(711\)](#), [Acceptance Table 4](#). Continue testing through all levels unless the results conform at an earlier level.

Test 2: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 2*.

Solution A: pH 6.4 phosphate buffer (21.7 g/L of [monobasic potassium phosphate](#) and 0.8 g/L of [sodium hydroxide](#) in [water](#), adjusted with 5 N [sodium hydroxide](#) or [phosphoric acid](#) to a pH of 6.4)

Solution B: 3.3 N [sodium hydroxide](#) (136 g/L of [sodium hydroxide](#) in [water](#))

Medium

Acid stage: 750 mL of 0.1 N [hydrochloric acid](#)

Buffer stage 1: 950 mL of *Solution A*

Buffer stage 2: 960 mL of pH 7.2 phosphate buffer

Apparatus 2: 100 rpm

Times

Acid stage: 2 h

Buffer stage 1: 1 h

Buffer stage 2: 1, 2, and 6 h

Acid stage

After 2 h of operation, withdraw a portion of the solution under test, discard the remaining solution, and retain the Tablets in proper order so that each will be returned later to its respective vessel. Blot the Tablets with a paper towel to dry and proceed immediately as directed in *Buffer stage 1*.

Standard solution: 0.016 mg/mL of [USP Mesalamine RS](#) in *Medium*. Sonicate to dissolve.

Sample solution: Pass a portion of the solution under test through a suitable filter of 0.45- μm pore size and discard the first few milliliters.

Instrumental conditions

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

Mode: UV

Analytical wavelength: 302 nm

Blank: *Medium*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of mesalamine ($\text{C}_7\text{H}_7\text{NO}_3$) dissolved:

$$\text{Result} = (A_U/A_S) \times C_S \times V \times (1/L) \times 100$$

A_U = absorbance of the *Sample solution*

A_S = absorbance of the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 750 mL

L = label claim of mesalamine (mg/Tablet)

Tolerances: NMT 1% of the labeled amount of mesalamine ($\text{C}_7\text{H}_7\text{NO}_3$) is dissolved.

Buffer stage 1

[NOTE—Use *Solution A* that has been equilibrated to a temperature of $37 \pm 0.5^\circ$.]

Transfer *Solution A* to each of the dissolution vessels, and place each Tablet from the *Acid stage* into its respective vessel. After 1 h, withdraw a 10-mL aliquot and proceed immediately as directed in *Buffer stage 2*.

Standard solution: 0.0125 mg/mL of [USP Mesalamine RS](#) in *Medium*. Sonicate to dissolve.

Sample solution: Pass a portion of the withdrawn solution under test through a suitable filter of 0.45- μm pore size and discard the first few milliliters.

Instrumental conditions

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

Mode: UV

Analytical wavelength: 330 nm

Blank: *Medium*

Analysis: Proceed as directed in the *Analysis* at *Acid stage*, using the *Medium* for *Buffer stage 1*.

Tolerances: NMT 1% of the labeled amount of mesalamine ($\text{C}_7\text{H}_7\text{NO}_3$) is dissolved.

Buffer stage 2

To adjust the pH of 940 mL of *Solution A* to pH 7.2, transfer 20 mL of *Solution B* into each dissolution vessel from *Buffer stage 1* and start the dissolution immediately.

At the end of the specified time point, withdraw 10 mL of the solution under test from each dissolution vessel and replace with 10 mL of *Medium* for *Buffer stage 2*.

Standard solution: 0.0315 mg/mL of [USP Mesalamine RS](#) in *Medium*. Sonicate to dissolve.

Sample solution: Dilute 2.5 mL of the withdrawn solution under test with *Medium* to 100 mL. Pass through a suitable filter of 0.45- μ m pore size and discard the first few milliliters.

Instrumental conditions

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

Mode: UV

Analytical wavelength: 332 nm

Blank: *Medium*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the concentration (C_i) of mesalamine ($C_7H_7NO_3$) in the sample withdrawn from the vessel at each time point (i):

$$\text{Result} = (A_U/A_S) \times C_S \times D$$

A_U = absorbance of the *Sample solution*

A_S = absorbance of the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

D = dilution factor of the *Sample solution*, 40

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved at each time point i :

$$\text{Result}_1 = C_1 \times V \times (1/L) \times 100$$

$$\text{Result}_2 = [(C_2 \times V) + (C_1 \times V_S)] \times (1/L) \times 100$$

$$\text{Result}_3 = \{(C_3 \times V) + [(C_2 + C_1) \times V_S]\} \times (1/L) \times 100$$

C_i = concentration of mesalamine in the portion of sample withdrawn at time point i (mg/mL)

V = volume of the *Medium*, 960 mL

L = label claim (mg/Tablet)

V_S = volume of the solution under test withdrawn at each time point (i) during *Buffer stage 2*, 10 mL

Tolerances: See [Table 2](#).

Table 2

Time Point (<i>i</i>)	Time (h)	Amount Dissolved (%)
1	1	NMT 35
2	2	35–60
3	6	NLT 80

Test 3: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 3*.

Solution A: pH 6.4 phosphate buffer [6.8 g/L of [potassium phosphate, monobasic](#) and 0.53 g/L of [sodium hydroxide](#) in [water](#), adjusted with 1 N (or 5 N) sodium hydroxide solution or [phosphoric acid](#) to a pH of 6.4]

Solution B: pH 7.2 phosphate buffer [6.8 g/L of [potassium phosphate, monobasic](#) and 1.4 g/L of [sodium hydroxide](#) in [water](#), adjusted with 1 N (or 5 N) sodium hydroxide solution or [phosphoric acid](#) to a pH of 7.2]

Medium

Acid stage: 0.1 N [hydrochloric acid](#), 750 mL

Buffer stage 1: *Solution A*, 950 mL

Buffer stage 2: *Solution B*, 960 mL

Apparatus 2

Acid stage: 100 rpm

Buffer stage 1: 100 rpm

Buffer stage 2: 100 rpm

Times

Acid stage: 2 h

Buffer stage 1: 1 h

Buffer stage 2: 1 h and 2 h

Buffer: 5 g/L of [potassium phosphate, monobasic](#) in [water](#), adjusted with [phosphoric acid](#) to a pH of 2.0 ± 0.05

Mobile phase: [Acetonitrile](#) and *Buffer* (20:80)

Standard solution: 1.25 mg/mL of [USP Mesalamine RS](#) in *Solution B*. Sonicate to dissolve.

Sample solutions

Acid stage: Place 1 Tablet in each vessel containing *Medium, Acid stage*. At the specified *Times*, withdraw a portion of the solution under test using a suitable filter of 10-µm pore size. Centrifuge if necessary. Remove the Tablets from solution, dry the Tablets with a paper towel, and retain in the proper order. Proceed as directed in *Buffer stage 1*.

Buffer stage 1: Transfer each Tablet from *Acid stage* into the respective vessel containing *Medium, Buffer stage 1*. At the specified *Times*, withdraw a portion of the solution under test using a suitable filter of 10-µm pore size. Centrifuge if necessary. Remove the Tablets from solution, dry the Tablets with a paper towel, and retain in the proper order. Proceed as directed in *Buffer stage 2*.

Buffer stage 2: Transfer each Tablet from *Buffer stage 1* into the respective vessel containing *Medium, Buffer stage 2*. At the specified *Times*, withdraw a portion of the solution under test using a suitable filter of 10-µm pore size. Centrifuge if necessary.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 330 nm

Column: 4.6-mm × 15-cm; 5-µm packing [L9](#)

Column temperature: 30°

Flow rate: 1.2 mL/min

Injection volume: 5 µL

Run time: NLT 2.5 times the retention time of mesalamine

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 2.0%

Analysis

Acid stage

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved:

$$\text{Result} = (r_U/r_S) \times C_S \times V \times (1/L) \times 100$$

r_U = peak response of mesalamine from the *Sample solution*

r_S = peak response of mesalamine from the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 750 mL

L = label claim (mg/Tablet)

Buffer stage 1: Proceed as directed for the *Acid stage* except the volume of *Medium* is 950 mL.

Buffer stage 2

Samples: *Standard solution* and *Sample solution*

Calculate the concentration (C_i) of mesalamine ($C_7H_7NO_3$) in the sample withdrawn from the vessel at each time point (i) as shown in [Table 3](#):

$$\text{Result}_i = (r_U/r_S) \times C_S$$

r_U = peak response of mesalamine from the *Sample solution*

r_S = peak response of mesalamine from the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved at each time point (i) as shown in [Table 3](#):

$$\text{Result}_1 = (C_1 \times V) \times (1/L) \times 100$$

$$\text{Result}_2 = \{[C_2 \times (V - V_S)] + (C_1 \times V_S)\} \times (1/L) \times 100$$

C_i = concentration of mesalamine in the portion of sample withdrawn at each time point (i) (mg/mL)

V = volume of *Medium*, 960 mL

L = label claim (mg/Tablet)

V_S = volume of the *Sample solution* withdrawn at each time point (i) (mL)

Tolerances

Acid stage: See [Table 3](#).

Buffer stage 1: See [Table 3](#).

Table 3

--

Level	Number Tested	Acceptance Criteria
L ₁	6	No individual value exceeds 1% dissolved.
L ₂	6	Average of the 12 units (L ₁ + L ₂) is NMT 1% dissolved, and no individual unit is >10% dissolved.
L ₃	12	Average of the 24 units (L ₁ + L ₂ + L ₃) is NMT 1% dissolved, and NMT 1 individual unit is >10% dissolved.

Buffer stage 2: See [Table 4](#).

Table 4

Level	Number Tested	Acceptance Criteria	
		Time Point 1 (1 h)	Time Point 2 (2 h)
L ₁	6	No individual value exceeds 65% dissolved.	Each unit is NLT 85% dissolved.
L ₂	6	Average of the 12 units (L ₁ + L ₂) is NMT 65% dissolved, and no individual unit is >75% dissolved.	Average of the 12 units (L ₁ + L ₂) is NLT 85% dissolved, and no unit is <75% dissolved.
L ₃	12	Average of the 24 units (L ₁ + L ₂ + L ₃) is NMT 65% dissolved, NMT 2 units are >75%, and no unit is >85% dissolved.	Average of the 24 units (L ₁ + L ₂ + L ₃) is NLT 85% dissolved, NMT 2 units are <75%, and no unit is <65% dissolved.

Test 4: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 4*.

Solution A: 0.2 M phosphate buffer (76.0 g/L of [sodium phosphate, tribasic, dodecahydrate](#) in [water](#))

Medium

Acid stage: 0.1 N [hydrochloric acid](#), 750 mL

Buffer stage 1: pH 6.4 phosphate buffer (after *Acid stage*, adjust with the addition of 200 mL of *Solution A* to the *Medium*, *Acid stage* to a pH of 6.4), 950 mL

Buffer stage 2: pH 7.2 phosphate buffer (after *Buffer stage 1*, adjust with the addition of about 10 mL of 2 M [sodium hydroxide](#) solution to the *Medium*, *Buffer stage 1* to a pH of 7.2), 960 mL

Apparatus 2: 100 rpm

Times

Acid stage: 2 h

Buffer stage 1: 1 h

Buffer stage 2: 1, 2, and 6 h

Acid stage

Standard solution: 0.016 mg/mL of [USP Mesalamine RS](#) in *Medium*. Sonicate to dissolve, if necessary.

Sample solution: At the specified *Times*, withdraw about 10 mL of the solution under test, pass through a suitable filter of 0.45-µm pore size, and collect the filtrate by discarding the first few milliliters.

Replace with about 10 mL of *Medium*.

Instrumental conditions

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

Mode: UV

Analytical wavelength: 302 nm

Cell: 1 cm

Blank: *Medium*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved:

$$\text{Result} = (A_U/A_S) \times C_S \times V \times (1/L) \times 100$$

A_U = absorbance of the *Sample solution*

A_S = absorbance of the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 750 mL

L = label claim of mesalamine (mg/Tablet)

Tolerances: NMT 1% of the labeled amount of mesalamine ($C_7H_7NO_3$) is dissolved.

Buffer stage 1

Standard solution: 0.012 mg/mL of [USP Mesalamine RS](#) in *Medium*. Sonicate to dissolve, if necessary.

Sample solution: At the specified *Times*, withdraw about 10 mL of the solution under test, pass through a suitable filter of 0.45- μ m pore size, and collect the filtrate by discarding the first few milliliters.

Replace with about 10 mL of *Medium*.

Instrumental conditions

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

Mode: UV

Analytical wavelength: 331 nm

Cell: 1 cm

Blank: *Medium*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved:

$$\text{Result} = (A_U/A_S) \times C_S \times V \times (1/L) \times 100$$

A_U = absorbance of the *Sample solution*

A_S = absorbance of the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 950 mL

L = label claim of mesalamine (mg/Tablet)

Tolerances: NMT 1% of the labeled amount of mesalamine ($C_7H_7NO_3$) is dissolved.

Buffer stage 2

Standard solution: 0.025 mg/mL of [USP Mesalamine RS](#) in *Medium*. Sonicate to dissolve, if necessary.

Sample solution: At the specified *Times*, withdraw about 10 mL of the solution under test, pass through a suitable filter of 0.45- μ m pore size, and collect the filtrate by discarding the first few milliliters. Dilute 2.0 mL of the filtrate with *Medium* to 100 mL. Replace with about 10 mL of *Medium*.

Instrumental conditions

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

Mode: UV

Analytical wavelength: 331 nm

Cell: 1 cm

Blank: *Medium*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the concentration (C_i) of mesalamine ($C_7H_7NO_3$) in the sample withdrawn from the vessel at each time point (i):

$$\text{Result} = (A_U/A_S) \times C_S \times D$$

A_U = absorbance of the *Sample solution*

A_S = absorbance of the *Standard solution*

C_S = concentration of [USP Mesalamine RS](#) in the *Standard solution* (mg/mL)

D = dilution factor of the *Sample solution*, 50

Calculate the percentage of the labeled amount of mesalamine ($C_7H_7NO_3$) dissolved at each time point (i):

$$\text{Result}_1 = C_i \times V \times (1/L) \times 100$$

$$\text{Result}_2 = [(C_2 \times V) + (C_1 \times V_S)] \times (1/L) \times 100$$

$$\text{Result}_3 = \{(C_3 \times V) + [(C_2 + C_1) \times V_S]\} \times (1/L) \times 100$$

C_i = concentration of mesalamine in the portion of sample withdrawn at each time point (i) (mg/mL)

V = volume of *Medium*, 960 mL

L = label claim (mg/Tablet)

V_S = volume of the solution under test withdrawn at each time point (i) during *Buffer stage 2*, 10 mL

Tolerances: See [Table 5](#).

Table 5

Time Point (<i>i</i>)	Time (h)	Amount Dissolved (%)
1	1	NMT 30
2	2	20–60
3	6	NLT 80

▲ (RB 1-Nov-2020)

- **UNIFORMITY OF DOSAGE UNITS** (905), *Weight Variation*: Meet the requirements

IMPURITIES

Change to read:

- **ORGANIC IMPURITIES**

Mobile phase, System suitability stock solution, System suitability solution, Standard stock solution, Standard solution, Chromatographic system, and System suitability: Proceed as directed in the Assay.

Sample solution: ▲ Prepare as directed for *Sample stock solution* in the Assay. ▲ (USP 1-Aug-2020)

Analysis

Sample: *Sample solution*

Calculate the percentage of each individual impurity in the portion of Tablets taken:

$$\text{Result} = (r_U/r_T) \times 100$$

r_U = peak response of each individual impurity

r_T = sum of all the peak responses

Acceptance criteria

Individual impurity: The largest secondary peak is NMT 1.0% of the total area.

Any other individual impurity: NMT 0.5%

Total impurities: NMT 2.0%

ADDITIONAL REQUIREMENTS

Change to read:

- **PACKAGING AND STORAGE:** Preserve in tight containers. ▲ Store at controlled room temperature. ▲ (USP 1-Aug-2020)

● **LABELING:** When more than one *Dissolution* test is given, the labeling states the *Dissolution* test used only if *Test 1* is not used.

- **USP REFERENCE STANDARDS** (11)

[USP Mesalamine RS](#)

[USP Salicylic Acid RS](#)

Page Information:

Not Applicable

DocID:

© 2020 The United States Pharmacopeial Convention *All Rights Reserved.*