

[x](#)

## How to Use

- **Searching:** Type keyword in search field at top of page. Search by all or part of a monograph title. For searches using multiple criteria, you will find items that match each of the specified criteria unless quotation marks are used.
  - For example, a search on Aminosalicyclic Acid Tablets will result in anything that contains “Aminosalicyclic” OR “Acid” OR “Tablets”
  - A search for “Aminosalicyclic Acid Tablets” will result in anything that specifically contains “Aminosalicyclic Acid Tablets”
- **Sorting:** Click on any column header title to sort alphabetically or chronologically in ascending or descending order. Note: the page load column is sorted alphabetically so that a number is ordered by first digit vs. by the actual number; thus, numbers will not always be in order.
  - For example, page 2178 will come before page 74 on a page sort.
- **Downloading:** You can download the Errata table in Comma-separated Value (.csv). The download will include the Errata that you have filtered on.
- **Importing:** You will need to import the file into Excel or Open Office with UTF-8 encoding, as opposed to simply opening it. To import, open Excel or Open Office and select import from the File drop-down. Depending on the version you are using, you should be presented with import formatting options to include UTF-8 as one of the first steps. Importing via UTF-8 should eliminate odd character conversions.

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
POWDERED	IDENTIFICATIO	<i>First</i>	5883	26-Jul-2013	1-Aug-2013	<i>USP38–NF33</i>	<i>First</i>	Line 2 of

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
GYMNEMA	N/B. Thin-Layer Supplement to Chromatograph USP36–NF31 y/ Chromatographic system							Supplement to USP37–NF32	Adsorbent: Change 5m to: 5 µm
GLYCERYL DISTEARATE	ASSAY/ Procedure/ Chromatographic system	USP36–NF31	2029	26-Jul-2013		1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	Line 1 of Column temperature: Change Column temperature: 40° to: Temperatures Detector: 40° Column: 40°
OXCARBAZEPINE	IMPURITIES/Organic Impurities, Procedure 2	First Supplement to USP36–NF31	6035	26-Jul-2013		1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	Row 9 of Column 1 of Table 3: Change Oxcarbazepine related compound E <sup>9</sup> to: Oxcarbazepine related compound E AND Delete footnote g
CEFDINIR	IM	USP36–NF31	2850	26-Jul-2013		1-Aug-2013	USP38–NF33	First	Row 16 of

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
CAPSULES	PURITIES/ <i>Organic Impurities/ Table 2</i>						<i>Supplement to USP37–NF32</i>	Column 1: Change Cefdinir impurity 2 <sup>e</sup> to: Cefdinir impurity 2 <sup>f</sup> AND Row 21 of Column 1: Change Cefdinir impurity 3 <sup>e</sup> to: Cefdinir impurity 3 <sup>f</sup>
DIPHENHYDRAMINE HYDROCHLORIDE CAPSULES	Assay	<i>USP36–NF31</i>	3276	26-Jul-2013	1-Aug-2013	<i>USP38–NF33</i>	<i>First Supplement to USP37–NF32</i>	Line 2: Change <i>Mobile phase, Standard preparation, System suitability solution, and Chromatographic system</i> —Prepare as directed in the Assay under <i>Diphenhydramine Hydrochloride</i> .

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>to:</p> <p><i>Mobile phase</i>—Prepare a solution of acetonitrile, water, and triethylamine (50: 50: 0.5), adjust with glacial acetic acid to a pH of 6.5, filter, and degas. Make adjustments if necessary (see <i>System Suitability</i> under <i>Chromatography</i> &lt;621&gt; ).</p> <p><i>Standard preparation</i></p> <p>—Dissolve an accurately weighed quantity of USP Diphenhydramine Hydrochloride RS in water to obtain a</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>solution having a known concentration of about 0.5 mg per mL.</p> <p>AND</p> <p>After the <i>Assay preparation</i> subsection: Add <i>System suitability solution</i></p> <p>—Dissolve about 5 mg of benzophenone in 5 mL of acetonitrile, dilute with water to 100 mL, and mix. Transfer 1.0 mL of this solution and 5 mg of diphenhydramine hydrochloride to a 10-mL volumetric flask, dilute with water to volume, and mix.</p> <p><i>Chromatographi</i></p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p><i>c system (see Chromatography &lt;621&gt;)</i>—The liquid chromatograph is equipped with a 254-nm detector and a 4.6-mm × 25-cm column that contains packing L10. The flow rate is about 1 mL per minute. Chromatograph the <i>System suitability solution</i>, and record the peak responses as directed for <i>Procedure</i>; the resolution, <i>R</i>, between the benzophenone and diphenhydramine peaks is not less than 2.0. Chromatograph</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>replicate injections of the <i>Standard preparation</i>, and record the peak responses as directed for <i>Procedure</i>; the relative standard deviation is not more than 2.0%, and the tailing factor for the diphenhydramine hydrochloride peak is not more than 2.0.</p> <p>AND</p> <p>Line 1 of <i>Procedure</i>: Change Proceed as directed for <i>Procedure</i> in the <i>Assay</i> under <i>Diphenhydramine Hydrochloride</i>. to:</p>

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
FELBAMATE TABLETS	IM PURITIES/ <i>Organic Impurities/System suitability/Suitability requirements</i>	USP36–NF31	3537	26-Jul-2013		1-Aug-2013	USP38–NF33	<i>First Supplement to USP37–NF32</i>	Separately inject equal volumes, about 10 µL, of the <i>Standard preparation</i> and the <i>Assay preparation</i> into the chromatograph, record the chromatograms, and measure the responses for the major peaks. Line 1 of <i>Resolution</i> : Change NMT 2 to: NLT 2
MOXIFLOXACIN OPTHALMIC SOLUTION	<i>Related compounds</i>	USP36–NF31	4414	26-Jul-2013		1-Aug-2013	USP38–NF33	<i>First Supplement to USP37–NF32</i>	Row 1 of Column 3 of <i>Table 2</i> : Change Relative Retention Time vs. Moxifloxacin



<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
VENLAFAXINE ADDITIONAL R TABLETS EQUIREMENT S/USP Reference Standards <11>	USP36–NF31	5554	26-Jul-2013	1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	to: Relative Retention vs. Moxifloxacin AND Row 1 of Column 3 of <i>Table 3</i> : Change Relative Retention Time vs. Moxifloxacin to: Relative Retention vs. Moxifloxacin Line 2 of Venlafaxine Related Compound A RS: Change 1-(1-(4-methoxy phenyl)-2-(meth ylamino)ethyl)cy clohexanol. C <sub>16</sub> H <sub>25</sub> NO <sub>2</sub> 263.38 to: 1-(1-(4-Methox yphenyl)-2-(met hylamino)ethyl)

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
POWDERED STINGING NETTLE	COMPOSITION /Content of Total Amino Acids	USP36–NF31	1606	26-Jul-2013		1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	cyclohexanol hydrochloride. $C_{16}H_{25}NO_2 \cdot HCl$ 299.84 Line 1 of Reagent solution: Change Solution containing 1.00 g of ninhydrin, 1.50 g of hydrindantin, to: Solution containing 1.00 g of ninhydrin, 150 mg of hydrindantin, Line 4 of Standard solution: Change Dissolve in 30% of the final flask volume, and dilute with water to volume. to: Dissolve in
MEPROBAMATE TABLETS	ASSAY/ Procedure	First Supplement to USP36–NF31	6015	26-Jul-2013		1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
BETAMETHAS ONE SODIUM PHOSPHATE	Identification/B. Thin-Layer Chromatographic Identification Test <201>	USP36–NF31	2645	26-Jul-2013		1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	30% of the final flask volume of acetonitrile, and dilute with water to volume. Line 1 of Test solution: Change 1 mg per mL to: 1 mg per mL in methanol.
DACARBAZINE FOR INJECTION	USP Reference standards <11>	USP36–NF31	3137	26-Jul-2013		1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	Line 3 of USP Dacarbazine Related Compound B RS: Change $C_4H_3N_5O$ 137.10 to: $C_4H_3N_5O \cdot H_2O$ 155.12
EDETATE DISODIUM	ASSAY/ Procedure	USP36–NF31	3370	26-Jul-2013		1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	Line 5: Delete Titrimetric system (See Titrimetry <541>.) Mode: Direct titration Titrant: 0.1 N sodium

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
LAMOTRIGINE PERFORMANC TABLETS E TESTS/ <i>Dissolution</i> <711>/Test 1	USP36–NF31	4056	26-Jul-2013	1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	hydroxide VS <i>Endpoint detection:</i> Visual Line 3 of <i>Standard solution:</i> Change 0.028 µg/mL to: 0.028 mg/mL Footnote c: Change 1-(3-Methoxyphenyl)-2-(dimethylaminomethyl)cyclohex-1-ene hydrochloride (identified and reported as an individual unspecified impurity if present). to: 1-(3-Methoxyphenyl)-2-(dimethylaminomethyl)cyclohex-6-ene hydrochloride
TRAMADOL HYIM DROCHLORID PUR E EXTENDED- RELEASE TABLETS	USP36–NF31	5438	26-Jul-2013	1-Aug-2013	USP38–NF33	First Supplement to USP37–NF32	ITIES/ <i>Organic Impurities/ Table</i> 2

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>(identified and reported as an individual unspecified impurity if present).  AND  Footnote d:  Change  1-(3-Methoxyphenyl)-2-(dimethylaminomethyl)cyclohex-6-ene hydrochloride  (identified and reported as an individual unspecified impurity if present).  to:  1-(3-Methoxyphenyl)-2-(dimethylaminomethyl)cyclohex-1-ene hydrochloride  (identified and reported as an individual unspecified</p>

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
PURIFIED GYMNEMA EXTRACT	COMPOSITION <i>/Content of Gymnemic Acids</i>	<i>First Supplement to USP36–NF31</i>	5884	26-Jul-2013		1-Aug-2013	<i>USP38–NF33</i>	<i>First Supplement to USP37–NF32</i>	impurity if present). Line 1 of <i>Acceptance criteria</i> : Change 90%–110% of the labeled amount to: 90.0%–110.0% of the labeled amount on the dried basis
SODIUM HYDROXIDE	ASSAY/ <i>Procedure</i>	<i>USP36–NF31</i>	2203	26-Jul-2013		1-Aug-2013	<i>USP38–NF33</i>	<i>First Supplement to USP37–NF32</i>	Line 11 of <i>Analysis</i> : Change = volume of <i>Titrant</i> consumed by the <i>Sample</i> to the first endpoint (mL) to: = volume of <i>Titrant</i> consumed by the <i>Sample</i> to the second endpoint (mL)
CLARITHROM YCIN FOR	ASSAY/ <i>Procedure</i>	<i>USP36–NF31</i>	3018	26-Jul-2013		1-Aug-2013	<i>USP38–NF33</i>	<i>First Supplement to</i>	Change the subsection

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
ORAL SUSPENSION						USP37–NF32	head <i>Buffer:</i> to: <i>Buffer A:</i> AND After the <i>Buffer A</i> subsection: Add <i>Buffer B:</i> 0.067 M dibasic potassium phosphate AND Line 1 of <i>Mobile phase:</i> Change Methanol and <i>Buffer</i> to: Methanol and <i>Buffer A</i> AND Line 4 of <i>Sample stock solution:</i> Change with the aid of 330 mL of <i>Buffer</i> , to a 1000-mL volumetric flask containing 50

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
PENTAZOCINE <i>Chemical Information</i> INJECTION	USP36–NF31	4734	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	mL of <i>Buffer</i> . to: with the aid of 330 mL of <i>Buffer B</i> , to a 1000-mL volumetric flask containing 50 mL of <i>Buffer B</i> . Line 1: Remove all chemical information.
POTASSIUM <i>Assay</i> CHLORIDE EX TENDED- RELEASE CAPSULES	USP36–NF31	4838	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	Line 2: Change <i>Potassium</i> <i>stock</i> <i>solution</i> and <i>Standard</i> <i>preparations</i> — to: <i>Standard stock</i> <i>solution</i> and <i>Standard</i> <i>solutions</i> — AND Line 1 of <i>Procedure</i> : Change for <i>Procedure</i> in the <i>Assay</i> under <i>Potassium</i> <i>Chloride Oral</i>



<a href="#">Monograph TitleSection</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
SELEGILINE H <i>Dissolution</i> YDROCHLORI <711> DE TABLETS	USP36–NF31	5120	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p><i>Solution.</i> to: for <i>Instrumental conditions</i> and <i>Analysis</i> in the <i>Assay</i> under <i>Potassium Chloride Oral Solution</i>, except use <i>Assay preparation</i> instead of <i>Sample solution</i>.</p> <p>Line 3 of <i>Chromatographic system</i>: Change <i>Chromatograph the Standard solution</i>, and record the peak responses. to: The flow rate is 1.0 mL/min. <i>Chromatograph the Standard solution</i>, and record the peak responses.</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
BIOTECHNOL METHODODOLOG OGY-DERIVED IES OF AMINO ARTICLES—AMACID NO ACID ANALYSIS ANALYSIS GENERAL PRINCIPLES	USP36–NF31	619	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	Change the section title <i>Method 6—Postcolumn DABS-Cl Derivatization General Principle</i> to: <i>Method 6—Precolumn DABS-Cl Derivatization General Principle</i>
VERAPAMIL H Assay YDROCHLORI DE ORAL SUSPENSION	USP36–NF31	5558	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	Line 2 of <i>Mobile phase</i> : Change 0.01 M to: 0.01 N AND Line 7 of <i>Assay preparation</i> : Change 10-mL to: 100-mL
AMITRIPTYLIN IDENTIFICATIO E HYDROCHL N/A. ORIDE TABLETS	USP36–NF31	2464	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	Line 2: Change <i>Sample solution</i> : Nominally 0.01

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>mg/mL of amitriptyline hydrochloride in methanol from a suitable amount of finely powdered Tablets. Filter a portion of the solution, and use the filtrate for analysis.</p> <p>to:</p> <p><i>Sample stock solution:</i> Nominally 0.1 mg/mL of amitriptyline hydrochloride in methanol from a suitable amount of finely powdered Tablets. Filter a portion of the solution, and use the filtrate.</p> <p><i>Sample solution:</i> Nominally 0.01 mg/mL of</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
MINOCYCLINE Assay FOR INJECTION	USP36–NF31	4375	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p>amitriptyline hydrochloride from <i>Sample stock solution</i> in methanol</p> <p>Line 2: Change <i>Mobile phase, Standard preparation, Resolution solution, and Chromatographic system</i>—Proceed as directed in the Assay under <i>Minocycline Hydrochloride</i>. to: <i>Mobile phase</i>—Prepare a mixture of 0.2 M ammonium oxalate, 0.01 M edetate disodium, dimethylformamide, and tetrahydrofuran (600:180:120:8</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>0). Adjust with ammonium hydroxide to a pH of 7.2, and pass through a filter of 0.5-µm or finer porosity. Make adjustments if necessary (see <i>System Suitability</i> under <i>Chromatography</i> &lt;621&gt;). <i>Standard preparation</i>—Dissolve an accurately weighed quantity of USP Minocycline Hydrochloride RS in water to obtain a solution having a known concentration of about 500 µg of minocycline (C</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							$^{23}\text{H}_{27}\text{N}_3\text{O}_7$ ) per mL. Use this solution within 3 hours. <i>Resolution solution</i> —Transfer 10 mg of USP Minocycline Hydrochloride RS to a 25-mL volumetric flask, add 20 mL of 0.2 M ammonium oxalate, and swirl to dissolve. Heat on a water bath at 60° for 180 minutes, and allow to cool. Dilute with water to volume, and mix. <i>Chromatographic system (see Chromatography &lt;621&gt;)</i> —The liquid

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>chromatograph is equipped with a 280-nm detector and a 4.6-mm x 25-cm column that contains 5-<math>\mu</math>m packing L1, and is maintained at a constant temperature of about 40°. The flow rate is about 1.5 mL per minute. Chromatograph the <i>Standard preparation</i>, and record the peak responses as directed for <i>Procedure</i>: the capacity factor, <math>k'</math>, is not less than 5.0 and not more than 11.5; the tailing factor for the analyte peak is not less</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>than 0.9 and not more than 2.0; and the relative standard deviation for replicate injections is not more than 2.0%.</p> <p>Chromatograph the <i>Resolution solution</i>, and record the peak responses as directed for <i>Procedure</i>: the relative retention times are about 0.7 for epiminocycline and 1.0 for minocycline; and the resolution, <i>R</i>, between epiminocycline and minocycline is not less than 4.6.</p> <p>AND</p>



<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
OXYCODONE AND ACETAMINOPHEN TABLETS	IDENTIFICATION N/A. <i>Thin-Layer Chromatography</i>	USP36–NF31	4645	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	<p>Line 1 of <i>Procedure</i>: Change Proceed as directed for <i>Procedure</i> in the Assay under <i>Minocycline Hydrochloride</i>. to: Separately inject equal volumes (about 20 µL) of the <i>Standard preparation</i> and the <i>Assay preparation</i> into the chromatograph, record the chromatograms, and measure the responses for the major peaks.</p> <p>Line 2 of <i>Sample solution</i>: Change in a mixture of</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
POTASSIUM Assay for BICARBONATE <i>potassium</i> AND POTASSIUM CHLORIDE EF FERVESCENT TABLETS FOR ORAL SOLUTION	USP36–NF31	4834	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	methanol and water (4:1). to: in a 5-mL mixture of methanol and water (4:1). Line 2: Change <i>Potassium stock solution</i> and <i>Standard preparations</i> — to: <i>Standard stock solution</i> and <i>Standard solutions</i> — AND Line 1 of <i>Procedure</i> : Change for <i>Procedure</i> in the Assay under <i>Potassium Chloride Oral Solution</i> . to: for <i>Instrumental conditions</i> and <i>Analysis</i> in the

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
POTASSIUM Assay CHLORIDE EX TENDED- RELEASE TABLETS	USP36–NF31	4841	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p>Assay under <i>Potassium Chloride Oral Solution</i>, except use Assay preparation instead of <i>Sample solution</i>.</p> <p>Line 5: Change <i>Potassium stock solution</i> and <i>Standard preparations</i>—to: <i>Standard stock solution</i> and <i>Standard solutions</i>—</p> <p>Line 1 of <i>Procedure</i>: Change in the Assay under <i>Potassium Chloride Oral Solution</i>. to: for <i>Instrumental conditions</i> and</p>

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
ISOTRETINOIN PERFORMANC CAPSULES	E TESTS/ Dissolution <711>/Test 4	Revision Bulletin (Official October 01, 2012)	Online	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	Analysis in the Assay under Potassium Chloride Oral Solution, except use Assay preparation 1 or Assay preparation 2 instead of Sample solution. Line 2 of Medium: Change 4.5% (v/v) of Milloxid L (lauryl dimethyl amine oxide) to: 4.5% (v/v) of lauryl dimethyl amine oxide
TROLAMINE SALICYLATE	Assay	USP36–NF31	5499	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	Line 3 of Chromatographic system: Change L1 to: L7
NITRIC ACID	ASSAY/	USP36–NF31	2107	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	Line 1 of

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
<i>Procedure</i>							<p><i>Sample solution:</i> Change To 2 mL of Nitric Acid in a tared, glass-stoppered conical flask add 25 mL of water. to: Weigh 2 mL of Nitric Acid in a glass-stoppered conical flask, and add 25 mL of water.</p>
FOSPHENYTOI Assay N SODIUM INJECTION	USP36–NF31	3680	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p>Line 1 of Assay preparation: Change Transfer an accurately measured volume of the Injection, equivalent to about 300 mg of fosphenytoin, to: Transfer an accurately</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
MOXIFLOXACIN OPHTHALMIC SOLUTION	USP36–NF31	4414	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	measured volume of the Injection, equivalent to about 300 mg of fosphenytoin sodium, Line 4 of <i>Resolution solution</i> : Change 0.1 mg per mg and 0.001 mg per mg, to: 0.1 mg per mL and 0.001 mg per mL,
POTASSIUM BICARBONATE EFFERVESCENT TABLETS FOR ORAL SOLUTION	USP36–NF31	4833	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	Line 2: Change <i>Potassium stock solution</i> and <i>Standard preparations</i> —to: <i>Standard stock solution</i> and <i>Standard solutions</i> —AND Line 1 of

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
POTASSIUM CHLORIDE FOR ORAL SOLUTION	Assay	USP36–NF31	4840	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	<p><i>Procedure:</i> Change for <i>Procedure</i> in the Assay under <i>Potassium Chloride Oral Solution</i>. to: for <i>Instrumental conditions</i> and <i>Analysis</i> in the Assay under <i>Potassium Chloride Oral Solution</i>, except use Assay preparation instead of <i>Sample solution</i>. Line 2: Change <i>Potassium stock solution</i> and <i>Standard preparations</i>— to: <i>Standard stock solution</i> and <i>Standard solutions</i>—</p>

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
TACROLIMUS CAPSULES	ADDITIONAL REQUIREMENT S/USP Reference Standards <11>	USP36–NF31	5257	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	<p>AND</p> <p>Line 1 of <i>Procedure</i>: Change for <i>Procedure</i> in the Assay under <i>Potassium Chloride Oral Solution</i>. to: for <i>Instrumental conditions</i> and <i>Analysis</i> in the Assay under <i>Potassium Chloride Oral Solution</i>, except use <i>Assay preparation 1</i> or <i>Assay preparation 2</i> instead of <i>Sample solution</i>.</p> <p>Line 11 of USP Tacrolimus System Suitability Mixture RS: Change and tacrolimus</p>



<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							8-propyl analog (3S,4R,5S,8S,9E,12S,14S,15R,16S,18R,19R,26aS)-5,6,8,11,12,13,14,15,16,17,18,19,24,25,26,26a-hexadecahydro-5,19-dihydroxy-3-((E)-2-[(1R,3R,4R)-4-hydroxy-3-methoxycyclohexyl]-1-methylvinyl)-14,16-dimethoxy-4,10,12,18-tetramethyl-15,19-epoxy-8-propyl-3H-pyridolo[2,1-c][1,4]oxaazacyclotricosine-1,7,20,21-(4H,23H)-tetrone. to: and tacrolimus 8-propyl analog (3S,4R,5S,8R

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
BIOTECHNOLOGY-DERIVED ARTICLES—AMINO ACID ANALYSIS	USP36–NF31	619	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	,9E,12S,14S,15R,16S,18R,19R,26aS)-5,6,8,11,12,13,14,15,16,17,18,19,24,25,26,26a-hexadecahydro-5,19-dihydroxy-3-((E)-2-[(1R,3R,4R)-4-hydroxy-3-methoxycyclohexyl]-1-methylvinyl)-14,16-dimethoxy-4,10,12,18-tetramethyl-15,19-epoxy-8-propyl-3H-pyridine[2,1-c][1,4]oxaazacyclotricosine-1,7,20,21-(4H,23H)-tetrone. Change the section title <i>Method 6—Postcolumn DABS-Cl Derivatization</i>

<a href="#">Monograph Title</a>	<a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a>	<a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
VITAMIN E	SPECIFIC TESTS/ <i>Acidity</i>	USP36–NF31	5579	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	to: <i>Method 6—Precolumn DABS-Cl Derivatization</i> Line 1 of <i>Sample:</i> Change 40 mg to: 1.0 g
FOSPHENYTOIN SODIUM	USP Reference standards <11>	USP36–NF31	3679	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	Line 6: Change C <sub>14</sub> H <sub>15</sub> NO <sub>2</sub> to: C <sub>14</sub> H <sub>13</sub> NO <sub>2</sub>
MINOCYCLINE HYDROCHLORIDE ORAL SUSPENSION	Assay	USP36–NF31	4376	31-May-2013		1-Jun-2013	USP37–NF32	USP37–NF32	Line 2: Change <i>Mobile phase</i> and <i>Chromatographic system</i> —Proceed as directed in the Assay under <i>Minocycline Hydrochloride</i> . to: <i>Mobile phase</i> —Prepare a mixture of 0.2 M ammonium

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>oxalate, 0.01 M edetate disodium, dimethylformamide, and tetrahydrofuran (600:180:120:80). Adjust with ammonium hydroxide to a pH of 7.2, and pass through a filter of 0.5-<math>\mu</math>m or finer pore size. Make adjustments if necessary (see <i>System Suitability</i> under <i>Chromatography</i> &lt;621&gt;). <i>Chromatographic system</i> (see <i>Chromatography</i> &lt;621&gt;)—The liquid chromatograph is equipped with a 280-nm detector and a 4.6-mm <math>\times</math></p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>25-cm column that contains 5-<math>\mu</math>m packing L1, and is maintained at a constant temperature of about 40°. The flow rate is about 1.5 mL per minute. Chromatograph the <i>Standard preparation</i>, and record the peak responses as directed for <i>Procedure</i>: the capacity factor, <math>k'</math>, is not less than 5.0 and not more than 11.5; the tailing factor for the analyte peak is not less than 0.9 and not more than 2.0; and the relative standard deviation for replicate</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>injections is not more than 2.0%.            Chromatograph the <i>Resolution solution</i>, and record the peak responses as directed for <i>Procedure</i>: the relative retention times are about 0.7 for epiminocycline and 1.0 for minocycline; and the resolution, <i>R</i>, between epiminocycline and minocycline is not less than 4.6.            AND            Line 1 of <i>Procedure</i>:            Change            Proceed as directed for <i>Procedure</i> in</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
OXYCODONE ASSAY/ TEREPHTHAL Procedure ATE	USP36–NF31	4648	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p>the Assay under <i>Minocycline Hydrochloride</i>. to: Separately inject equal volumes (about 20 µL) of the <i>Standard preparation</i> and the Assay <i>preparation</i> into the chromatograph, record the chromatograms, and measure the responses for the major peaks.</p> <p>Line 7 of <i>Analysis</i>: Change <math>R_U</math> = internal standard ratio (peak response of oxycodone/peak response of ethylparaben) from the <i>Standard</i></p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
POTASSIUM CHLORIDE EX <711> TENDED-RELEASE CAPSULES	USP36–NF31	4838	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p><i>solution</i>  <math>R_S</math> = internal standard ratio (peak response of oxycodone/peak response of ethylparaben) from the <i>Sample solution</i> to:  <math>R_U</math> = peak response ratio of oxycodone to ethylparaben from the <i>Sample solution</i>  <math>R_S</math> = peak response ratio of oxycodone to ethylparaben from the <i>Standard solution</i></p> <p>Line 5: Change <i>Potassium stock solution</i> and <i>Standard preparations</i>—to:  <i>Standard stock</i></p>



<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
POTASSIUM CHLORIDE, POTASSIUM BICARBONATE, AND POTASSIUM CITRATE EFFERVESCENT TABLETS FOR ORAL SOLUTION	<i>Assay for potassium</i> USP36–NF31	4843	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p><i>solution and Standard solutions—</i> AND Line 7 of <i>Procedure:</i> Change for <i>Procedure</i> in the Assay under <i>Potassium Chloride Oral Solution.</i> to: for <i>Instrumental conditions and Analysis</i> in the Assay under <i>Potassium Chloride Oral Solution.</i></p> <p>Line 2: Change <i>Potassium stock solution and Standard preparations—</i> to: <i>Standard stock solution and Standard solutions—</i></p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
ETHYLENE OXIDE AND DIOXANE	<i>Method II</i>	USP36–NF31 148	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p>AND</p> <p>Line 1 of <i>Procedure</i>: Change for <i>Procedure</i> in the Assay under <i>Potassium Chloride Oral Solution</i>. to: for <i>Instrumental conditions</i> and <i>Analysis</i> in the Assay under <i>Potassium Chloride Oral Solution</i>, except use <i>Assay preparation</i> instead of <i>Sample solution</i>.</p> <p>Line 31 of <i>Analysis</i>: Change <math>r_S</math> = ethylene oxide peak responses from <i>Standard solution B</i> to:</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
VERAPAMIL H YDROCHLORI DE ORAL SOLUTION	Assay <i>USP36–NF31</i>	5558	31-May-2013	1-Jun-2013	<i>USP37–NF32</i>	<i>USP37–NF32</i>	<p><math>r_s</math> = dioxane peak responses from <i>Standard solution B</i></p> <p>Line 3 of <i>Sodium acetate solution</i>: Change 0.01 M to: 0.01 N AND Line 6 of <i>Assay preparation</i>: Change 10-mL to: 100-mL</p>
ALBUTEROL SULFATE	<i>Chromatographic purity</i> <i>USP36–NF31</i>	2352	31-May-2013	1-Jun-2013	<i>USP37–NF32</i>	<i>USP37–NF32</i>	<p>Line 1: Change It meets the requirements of the test for <i>Chromatographic purity</i> under <i>Albuterol</i>, except to read Albuterol Sulfate in place of Albuterol and to use water instead of</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
MINOCYCLINE Assay HYDROCHLOR IDE CAPSULES	USP36–NF31	4375	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	<p>methanol as the solvent to prepare the <i>Standard solution</i> and the <i>Test solution</i>. to: It meets the requirements of the test for <i>Organic Impurities</i> under <i>Albuterol</i>, except to read Albuterol Sulfate in place of Albuterol and to use water instead of methanol as the solvent to prepare the <i>Standard solution</i> and the <i>Sample solution</i>.</p> <p>Line 2: Change <i>Mobile phase, Standard preparation, Resolution</i></p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p><i>solution, and Chromatographic system</i></p> <p>—Proceed as directed in the Assay under <i>Minocycline Hydrochloride</i>.</p> <p>to:</p> <p><i>Mobile phase</i>—Prepare a mixture of 0.2 M ammonium oxalate, 0.01 M edetate disodium, dimethylformamide, and tetrahydrofuran (600:180:120:80). Adjust with ammonium hydroxide to a pH of 7.2, and pass through a filter of 0.5-<math>\mu</math>m or finer pore size. Make adjustments if necessary (see</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p><i>System Suitability</i> under <i>Chromatography</i> &lt;621&gt;).  <i>Standard preparation</i>  —Dissolve an accurately weighed quantity of USP Minocycline Hydrochloride RS in water to obtain a solution having a known concentration of about 500 µg of minocycline (C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>7</sub>) per mL. Use this solution within 3 hours.</p> <p><i>Resolution solution</i>  —Transfer 10 mg of USP Minocycline Hydrochloride RS to a 25-mL</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>volumetric flask, add 20 mL of 0.2 M ammonium oxalate, and swirl to dissolve. Heat on a water bath at 60° for 180 minutes, and allow to cool. Dilute with water to volume, and mix.</p> <p><i>Chromatographic system (see Chromatography &lt;621&gt;)</i>—The liquid chromatograph is equipped with a 280-nm detector and a 4.6-mm x 25-cm column that contains 5-µm packing L1, and is maintained at a constant</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p>temperature of about 40°. The flow rate is about 1.5 mL per minute. Chromatograph the <i>Standard preparation</i>, and record the peak responses as directed for <i>Procedure</i>: the capacity factor, <math>k'</math>, is not less than 5.0 and not more than 11.5; the tailing factor for the analyte peak is not less than 0.9 and not more than 2.0; and the relative standard deviation for replicate injections is not more than 2.0%. Chromatograph the <i>Resolution</i></p>



<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
							<p><i>solution</i>, and record the peak responses as directed for <i>Procedure</i>: the relative retention times are about 0.7 for epiminocycline and 1.0 for minocycline; and the resolution, <i>R</i>, between epiminocycline and minocycline is not less than 4.6.</p> <p>AND</p> <p>Line 1 of <i>Procedure</i>: Change Proceed as directed for <i>Procedure</i> in the Assay under <i>Minocycline Hydrochloride</i>. to: Separately</p>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
NIFEDIPINE EXPERFORMANC TENDEE- RELEASE TABLETS	USP36–NF31	4509	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	inject equal volumes (about 20 µL) of the <i>Standard preparation</i> and the <i>Assay preparation</i> into the chromatograph, record the chromatograms, and measure the responses for the major peaks. Line 1 of <i>Solution A</i> : Change Dissolve 330.9 mg of sodium phosphate to: Dissolve 330.9 g of dibasic sodium phosphate Line 2: Change <i>Potassium stock solution</i> and <i>Standard</i>
POTASSIUM Assay for BICARBONATE <i>potassium</i> AND POTASSIUM CHLORIDE	USP36–NF31	4834	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
FOR EFFERVE SCENT ORAL SOLUTION							<p><i>preparations— to: Standard stock solution and Standard solutions— AND Line 1 of Procedure: Change for Procedure in the Assay under Potassium Chloride Oral Solution. to: for Instrumental conditions and Analysis in the Assay under Potassium Chloride Oral Solution, except use Assay preparation instead of Sample solution.</i></p>
POTASSIUM <i>Dissolution</i> CHLORIDE EX <711> TENDED-	USP36–NF31	4841	31-May-2013	1-Jun-2013	USP37–NF32	USP37–NF32	Line 5: Change <i>Potassium stock solution—</i>

<a href="#">Monograph Title</a> <a href="#">Section</a>	<a href="#">Source Publication</a>	<a href="#">Page Number</a>	<a href="#">Errata Post Date</a> <a href="#">Sort ascending</a>	<a href="#">Errata Official Date</a>	<a href="#">Target Errata Print Publication</a>	<a href="#">Target Online Fix Publication</a>	Description
RELEASE TABLETS							<p>to:  <i>Standard stock solution—</i>  AND  Line 7: Change  Prepare as directed for  <i>Standard preparations</i>  to:  Prepare as directed for  <i>Standard solutions</i>  AND  Line 7 of  <i>Procedure:</i>  Change  for <i>Procedure</i> in  the Assay under  <i>Potassium Chloride Oral Solution.</i>  to:  for <i>Instrumental conditions</i> and  <i>Analysis</i> in the  Assay under  <i>Potassium Chloride Oral Solution.</i></p>

---

## Pagination

- [\\_ First page « First](#)
- [\\_ Previous page ‹ Previous](#)
- ...
- [\\_ Page 32](#)
- [\\_ Page 33](#)
- [\\_ Page 34](#)
- [\\_ Page 35](#)
- [\\_ Page 36](#)
- [\\_ Page 37](#)
- [\\_ Page 38](#)
- [\\_ Page 39](#)
- [\\_ Page 40](#)
- [\\_ Next page Next ›](#)
- [\\_ Last page Last »](#)